
RESEARCH

BMC Rheumatology



Efficacy and safety of baricitinib in treatment of systemic lupus erythematosus: a systematic review and meta-analysis

Abdallah R. Allam^{1*}, Mohamed Salah Alhateem¹ and Abdelrahman Mohamed Mahmoud¹

Abstract

Background SLE is an autoimmune disease marked by broad immunological dysregulation and multi-system inflammation. Baricitinib is one of the novel treatments for SLE. We conducted this meta-analysis to evaluate its safety and effectiveness in treating SLE.

Method We looked for all published randomized controlled trials in PubMed, Scopus, Web of Science, and Cochrane and included all RCTs comparing baricitinib and placebo in the treatment of SLE. Review Manager 5.4 program was used for data analysis.

Results Three trials with a total of 1849 individuals were included. Participants in the baricitinib group were significantly more likely to attain SRI-4 response than those in the placebo group [RR = 1.11, 95% CI (1.02, 1.21), P=0.01]. Additionally, baricitinib performed better than the placebo in terms of reduction of \geq 4 points from baseline in SLEDAI-2 K score [RR = 1.13, 95% CI (1.04, 1.22), P=0.004]. In terms of SLEDAI-2 K remission of arthritis or rash, baricitinib was also superior to placebo [RR = 1.08, 95% CI (1.00, 1.17), P=0.04]. Treatment-emergent adverse events did not differ significantly [RR = 1.01, 95% CI (0.97, 1.05), P=0.61].

Conclusion Baricitinib is potentially safe and effective in the treatment of SLE. It has successfully met the study's primary endpoint and some secondary endpoints highlighting its potential to improve the outcomes of SLE. Despite achieving an SRI-4 response, glucocorticoids sparing and some other secondary outcomes weren't reached by baricitinib.

Keywords Baricitinib, Systemic Lupus Erythematosus, SLE, SRI-4, Systematic review, Meta-analysis

*Correspondence: Abdallah R. Allam abdallahreda89@med.menofia.edu.eg

¹Faculty of Medicine, Menoufia University, Yassin Abdelghaffar Street from Gamal Abdelnaser Street. Shebin Al- Kom, Menoufia 32511, Egypt



© The Author(s) 2023. **Open Access** This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit http://creativecommons.org/licenses/by/4.0/. The Creative Commons Dublic Domain Dedication waiver (http://creativecommons.org/publicdomain/zero/1.0/) applies to the data made available in this article, unless otherwise stated in a credit line to the data.

Introduction

Systemic lupus erythematosus (SLE) is a chronic autoimmune disease that affects numerous body organs and systems [1]. It is caused by the interplay between genetic, immunological, and environmental factors and presents with a wide range of symptoms like joint pain, skin rash, and multi-organ damage [2].

SLE has no known cause or treatment; nevertheless, there are several ways to manage the condition and slow its progression. These drugs include a combination of NSAIDs, anti-malarial drugs, corticosteroids, and immunosuppressive agents [3]. However, these medications are linked to substantial adverse events, and many patients suffer from disease relapse and symptom flare-ups [4].

The development of new treatments for SLE has been challenging, with few advancements in the past six decades. Only two new treatments approved in the last 60 years targeting B cells and type 1 interferon highlighting the unmet need for the development of new SLE therapies [5, 6].

The pathogenesis of SLE is complex, involving various immune mechanisms and factors underlying the disease activity. Different patients and clinical presentations exhibit variations in the dominance of these mechanisms. Consequently, the existing limited range of therapeutic targets does not cater to all patients. Extensive research has implicated a range of cytokines, including interferons, B-cell activating factors, various interleukins, and TNF in the development of SLE [7–9]. Many of these cytokines rely on Janus kinases (JAKs) for intracellular signaling [1, 10, 11].

Janus kinase inhibitors, a group of drugs approved by the US Food and Drug Administration, have shown an effective role in treating many inflammatory and autoimmune diseases, such as rheumatoid arthritis and psoriasis [3, 12]. They work by blocking the pathways of several cytokines, including interleukin (IL)-6 and interferon and interferon (IFN)- γ , which are involved in the pathogenesis of SLE [11, 13].

Baricitinib, one of Janus kinase inhibitors, has emerged as a potential therapeutic option for the treatment of a wide scale of autoimmune conditions including SLE [14– 17]. A number of clinical trials have been carried out to assess the safety and efficacy of baricitinib in the treatment of SLE [15–17]. However, the results of these trials were conflicting, Morand et al. reported potential effectiveness of baricitinib 4 mg in treatment of SLE which was supported by Wallace et al. On the other hand, Petri et al. reported failure of baricitinib to meet the primary end point of the study. Therefore, it is still uncertain if baricitinib is effective for SLE treatment.

We carried out this comprehensive systematic review to compare the safety and effectiveness of baricitinib versus placebo in the treatment of SLE. Moreover, we conducted a meta-analysis of available data to estimate the effect of baricitinib on disease activity, flare-ups, and adverse events.

Methods and materials

We performed this systematic review according to Cochrane guidelines [18] and preferred reporting items for systematic reviews and meta-analyses (PRISMA) guidelines [19].

Literature search and screening

We performed a systematic search on PubMed, Scopus, Web of Science, and Cochrane till March 2023. Entry terms were as follows: ("Systemic Lupus Erythematosus," "Lupus Erythematosus Disseminatus," "Libman-Sacks Disease," "Disease, Libman-Sacks," "Libman Sacks Disease," "baricitinib," "Olumiant," "baricitinib phosphate," "baricitinib phosphate salt."

In order to find pertinent publications for our systematic review, we carried out a four-step screening procedure. In the initial phase, we collected all retrieved articles and eliminated duplicates using Endnote 20 software. Next, the articles' titles and abstracts were inspected to rule out studies that weren't relevant. In the third step, we examined the complete texts of the remaining articles to determine whether they qualified for the review. Finally, we looked over the reference lists of the articles that were included to identify any potentially relevant publications that were missed in the initial search. The screening process was conducted independently by two researchers using the SR Accelerator tool, and any disagreements were resolved by a third researcher [20].

Eligibility criteria

We restricted our analysis to baricitinib versus placebo comparisons in randomized clinical trials for the treatment of SLE. Observational studies, case studies, conference abstracts, and in vitro and animal experiments were all disregarded.

Risk of bias assessment

We used the Cochrane risk of bias assessment tool (ROB2) to evaluate the quality of the included articles [21]. ROB2 compromises six domains: the randomized process, deviations from intended interventions, missing outcome data, measurement of outcome, choice of reported findings, and overall risk of bias.

Data extraction

Using a predetermined data extraction sheet, two different researchers independently extracted the data from the included studies, and any inconsistencies were resolved by a third researcher. Study ID, study design, NCT, site, length of treatment, inclusion criteria, exclusion criteria, and conclusion were included in the summary data. Age, time since onset of SLE, concomitant medications, baseline Systemic Lupus Erythematosus Disease Activity Index-2000 (SLEDAI-2 K) score, (SLEDAI-2 K) score organ system involvement, Physician's Global Assessment of Disease Activity (PGA), Cutaneous Lupus Erythematosus Disease Area and Severity Index (CLASI) activity score, tender joint counts (TJC), swollen joint counts (SJC), Systemic Lupus International Collaborating Clinics/ American College of Rheumatology (SLICC/ACR) Damage Index score was among the baseline characteristics.

Study outcomes

Efficacy outcomes included Systemic Lupus Erythematosus Responder Index-4 (SRI-4) as the primary outcome, reduction of 4 points or more from baseline in Systemic Lupus Erythematosus Disease Activity Index-2000 (SLE-DAI-2 K) score, SLEDAI-2 K remission of arthritis or rash, no new British Isles Lupus Assessment Group (BILAG) A and no more than one new (BILAG) B disease activity score, no worsening in Physician Global Assessment (PGA), glucocorticoid sparing, lupus low disease activity state (LLDAS). while safety outcomes included Treatment-emergent adverse events (TEAEs) Discontinuation from study treatment because of an adverse event, infections, serious infections, Opportunistic infections, hepatic disorders, and serious adverse effects.

Data analysis

Meta-analysis

In order to compare the baricitinib group with the placebo group, we used Review Manager version 5.4. To determine the differences in safety and efficacy between baricitinib 2 mg and baricitinib 4 mg, we performed subgroup analysis based on dose. For dichotomous data, the Mantel-Haenszel technique was used to display the risk ratio (RR) and 95% confidence intervals (CI); for continuous data, the inverse variance method was used to display the mean difference (MD) and 95% CI. When P < 0.05, there were significant differences. To assess the degree of heterogeneity, we conducted Chi-Square and I^2 tests. When the $I^2 > 50\%$ and the P-value of the Chisquare<0.1, there was significant heterogeneity. If the data were heterogeneous, we would apply a leave-oneout sensitivity analysis or a random effect model. A fixed effect model was utilized in the other cases.

Meta-regression

Using Comprehensive meta-analysis (CMA) software, we conducted meta regression analysis to determine if the dose of baricitinib, age of patients, sample size of the study and time since onset of SLE have influenced the primary outcome (SRI-4).

Results

After removing duplicates with the systematic review accelerator tool [20], the search turned up a total of 267 citations out of 376 citations. Additionally, two writers looked at the title and abstract of 267 citations and concluded that 16 of them could move on to the full-text screening stage. Finally, three of the 16 citations were included in this meta-analysis [15–17]. All discrepancies were resolved by the third author. PRISMA flow diagram is shown in. (Fig. 1)

In our study, there were 1849 participants, with a mean age of 43.1 years, a mean SLEDAI-2 K score of 9.7, and a prevalence of glucocorticoid use of 77.3%. 621 (33.6%) received 2 mg of baricitinib, 614 (33.2%) received 4 mg of baricitinib, and 614 (33.2%) received a placebo. The summary and baseline characteristics of the included studies are displayed in (Table 1) and (Table 2).

Risk of bias results

All of the included studies had a low risk of bias. (Fig. 2) and (S1) respectively exhibit a graph and summary of the bias risk.

Meta-analysis

Efficacy outcomes

SRI-4 Participants in the baricitinib group were significantly more likely to obtain an SRI-4 response than those in the placebo group [RR=1.13, 95% CI (1.04, 1.22), P=0.004]. Moreover, baricitinib 4 mg significantly outperformed placebo in the sub-group analysis [RR=1.18, 95% CI (1.06, 1.32), P=0.003]. (Fig. 3)

Reduction of \geq **4 points from baseline in SLEDAI-2 K score** Participants in the baricitinib group were significantly more likely to obtain a reduction of \geq 4 *points from baseline in SLEDAI-2 K score* than those in the placebo group [RR=1.13, 95% CI (1.04, 1.22), P=0.004]. Moreover, baricitinib 4 mg significantly outperformed the placebo in the sub-group analysis [RR=1.18, 95% CI (1.06, 1.32), P=0.003]. (Fig. 4)

SLEDAI-2 K remission of arthritis or rash Participants in the baricitinib group were significantly more likely to obtain SLEDAI-2 K remission of arthritis or rash than those in the placebo group [RR=1.08, 95% CI (1.00, 1.17), P=0.04]. Moreover, baricitinib 4 mg significantly outperformed placebo in the sub-group analysis [RR=1.14, 95% CI (1.03, 1.27), P=0.01]. (Fig. 5)




Fig. 1 PRISMA flow diagram

No new BILAG A and no more than one new BILAG B disease activity score In terms of BILAG, there was no significant difference between the baricitinib group and the placebo group [RR=1.03, 95% CI (0.98, 1.07), P=0.25]. Similarly, there was no significant difference between

baricitinib 4 mg and placebo in the sub-group analysis [RR=1.03, 95% CI (0.97, 1.10), P=0.32]. The data, however, were heterogeneous $[P=0.11, I^2=56\%]$. I took out Petri et al. [17] to address this heterogeneity. The outcome was altered and baricitinib 4 mg was significantly better

₽	NCT	Study design	Site	Dura- tion of TTT	Inclusion criteria	Exclusion criteria	Conclusion
Petri 2023	NCT03616964	Phase 3	USA, Central America, Mexico, Argentina, Chile, Colombia, France, Italy, Poland, Romania, Spain, Japan, Korea, Philip- pines, India, Serbia, and South Africa.	52 weeks	Participants required to be at least 18 years old, have a clinical diagnosis of SLE at least 24 weeks before screening, and meet four out of the eleven updated ACR criteria for SLE categorization. They also had a total SLEDAI-2 K score of at least 6 at screening and a clinical SLEDAI-2 K score of at least 4 at baseline, at least one BILAG A score or two BILAG B scores and were positive for at least one of the follow-ing: ANA, anti-dsDNA, or anti-Smith.	If a participant had severe active lupus nephritis, ac- tive CNS lupus, or another systemic inflammatory disease besides SLE, they were excluded from the study.	Baricitinib was proposed as a po- tential treatment for SLE patients in phase 2 data, which was sup- ported in SLE-BRAVE-I, however this finding was not repeated in SLE-BRAVE-II. There were no new warning signs of safety.
Mo- rand 2023	NCT03616912	Phase 3	China, Taiwan, Central America, Mexico, Brazil, Aus- tria, Belgium, Croatia, Czech Republic, Germany, Greece, Hungary, Switzerland, Netherlands, UK, USA, Australia, and Russia.	52 weeks	Participants required to be at least 18 years old, have a clinical diagno- sis of SLE at least 24 weeks before screening, and meet four out of the eleven updated ACR criteria for SLE categorization. Additionally, they had at least one ANA (titre \geq 1.80), anti-dsDNA, or anti-Smith positive screening result; active disease indicated by a screening SLEDAI-2 K score of at least 6; and at least one BILAG A score or two BILAG B scores despite SOC treatments.	Patients were disqualified if they had severe active CNS lupus, severe active lupus nephritis, had received treatment for, or were currently experienc- ing, another systemic inflammatory disease other than SLE.	The 4 mg baricitinib group in this study achieved the primary goal. However, important auxilia- ry endpoints weren't. There were no new warning signs of safety.
Wal- lace 2018	NCT02708095 Phase 2	Phase 2	11 countries in Asia, Europe, North America, and South America.	weeks	Participants were to be 18 years of age or older and have received a diagnosis of SLE at least 24 weeks prior to screening by meeting four or more of the 2012 SLICC classification criteria or the new ACR criteria for SLE classification. Patients had to have a positive anti-dSDNA; 2 30 IU/mL, a positive ANA test (HEp-2 titre 1:80), a positive SLEDAI-2 K score of 4 or above based on clinical symptoms, or active arthritis or rash as defined by the SLEDAI-2 K. The study medicine was added to already-present stable background SOC, which may have included corticosteroids or nonsteroidal anti-inflammatory drugs.	Active severe lupus ne- phritis, active severe CNS lupus, a recent clinically acute infection, and a few specific test abnormalities were important exclusion criteria.	With a safety profile consistent with prior baricitinib studies, the 4 mg dose of baricitinib, but not the 2 mg dose, significantly im- proved the signs and symptoms of active SLE in patients who were not sufficiently managed despite SOC. This research lays the groundwork for upcoming phase 3 trials using the possible oral treatment for SLE, baricitinib.

Allam et al. BMC Rheumatology (2023) 7:40




Fig. 2 Risk of bias graph

than the placebo [RR=1.09, 95% CI (1.01, 1.19), P=0.03]. (Fig. 6A&B)

No worsening by PGA In terms of PGA, there was no significant difference between the baricitinib group and the placebo group [RR=1.03, 95% CI (0.98, 1.07), P=0.25]. (Fig. 7)

Glucocorticoid sparing The glucocorticoid-sparing outcome was reported in two studies [16, 17]. There was no significant difference between the baricitinib group and the placebo group [RR=1.02, 95% CI (0.84, 1.24), P=0.86]. (Figure S2)

LLDAS In terms of LLDAS, there was no significant difference between the baricitinib group and the placebo group [RR=1.12, 95% CI (0.98, 1.27), P=0.1]. (Figure S3)

Participants with \geq **1 severe flare** In terms of participants with \geq 1 severe flare, there was no significant differ-

ence between the baricitinib group and the place bo group [RR=0.88, 95% CI (0.70, 1.09), P=0.24]. (Figure S4)

SLEDAI-2 K "score" The baricitinib group achieved a lower SLEDAI-2 K score than the placebo group, indicating a significant difference between the two groups [MD = -0.44, 95% CI (-0.72, -0.15), P=0.003]. Moreover, baricitinib 4 mg significantly outperformed the placebo in the sub-group analysis [MD = -0.67, 95% CI (-1.08, -0.26), P=0.001]. (Figure S5)

Safety outcomes

TEAEs The baricitinib group and the placebo group did not differ significantly in terms of TEAEs [RR=1.01, 95% CI (0.97, 1.05), P=0.61]. Moreover, there was no significant difference between the baricitinib group and the pla-

	Experim	ental	Contr	ol		Risk Ratio	Risk Ratio
Study or Subgroup	Events	Total	Events	Total	Weight	M-H, Fixed, 95% CI	M-H, Fixed, 95% Cl
1.1.1 2 mg							
Morand 2023	126	255	116	253	20.6%	1.08 [0.90, 1.29]	i +
Petri 2023	120	261	116	256	20.7%	1.01 [0.84, 1.22]	i +
Wallace 2018 Subtotal (95% CI)	54	105 621	50	105 614	8.8% 50.1%	1.08 [0.82, 1.42] 1.05 [0.93, 1.18]	
Total events	300	021	282	014	50.170	1.00 [0.00, 1110]	' T
Heterogeneity: Chi ² =	0.24, df =	2 (P = 0	.89); I ² =	0%			
Test for overall effect	and the second second	•					
1.1.2 4 mg							
Morand 2023	142	252	116	253	20.5%	1.23 [1.03, 1.46]	•
Petri 2023	121	258	116	256	20.6%	1.04 [0.86, 1.25]	
Wallace 2018	67	104	50	105	8.8%	1.35 [1.06, 1.73]	· · ·
Subtotal (95% CI)		614		614	49.9%	1.17 [1.05, 1.31]	I I I I I I I I I I I I I I I I I I I
Total events	330		282				
Heterogeneity: Chi ² =	: 3.30, df =	2 (P = 0	.19); I ^z =	39%			
Test for overall effect	: Z= 2.74 (P = 0.00	6)				
Total (95% CI)		1235		1228	100.0%	1.11 [1.02, 1.21]	•
Total events	630		564				
Heterogeneity: Chi ² =		•		7%			0.01 0.1 1 10 100
Test for overall effect			·	nst et	100 B	(margine to	Favours [control] Favours [experimental]
Test for subgroup dif	ferences: (Chi² = 1.	64. df = 1	(P = 0	.20), I ^z = 3	39.2%	

Fig. 3 A forest plot displaying the prevalence of SLE Responder Index-4 in the baricitinib group versus the placebo group

2	Duration	Dose	Age (years),	Time since	Concomital	Concomitant medications						
			M (SD)	onset of SLE, M (SD)	Glucocor- ticoids, N (%)	Prednisone dose (mg/ day), M (SD)	Antimalari- als, N (%)	Immunosup- pressants, N (%)	Metho- trexate, N (%)	Azathio- prine, N (%)	Mycopheno- late mofetil, N (%)	NSAIDs, N (%)
Petri 2023	52 weeks	2 mg	42.8 (13.0)	8.7 (7.7)	210 (80%)	9.6 (5.9)	213 (82%)	133 (51%)	52 (20%)	48 (18%)	25 (10%)	64 (25%)
		4 mg	42.2 (12.1)	8.5 (7.7)	207 (80%)	9.8 (5.7)	213 (83%)	133 (52%)	60 (23%)	33 (13%)	25 (10%)	70 (27%)
		Placebo	43.5 (13.5)	9-0 (8-3)	207 (81%)	8.8 (5.0)	209 (82%	140 (55%)	49 (19%)	47 (18%)	29 (11%)	51 (20%)
Morand 2023	52 weeks	2 mg	42.9 (12.4)	9.2 (7.7)	194 (76%)	10.4 (7)	189 (74%)	152 (60%)	51 (20%)	54 (21%)	39 (15%)	67 (26%)
		4 mg	41.5 (12.9)	8-8 (8-2)	187 (74%)	10.1 (6)	206 (82%)	141 (56%)	52 (21%)	42 (17%)	34 (14%)	68 (27%)
		Placebo	42.0 (12.0)	9.4 (7.5)	195 (77%)	9.8 (5)	213 (84%)	150 (59%)	60 (24%)	38 (15%)	39 (15%)	63 (25%)
Wallace 2018	24 weeks	2 mg	43.2 (11.0)	11-8 (9-1)	79 (75%)	8.7 (5.8)	71 (68%)	47 (45%)	17 (16%)	10 (10%)	10 (10%)	29 (28%)
		4 mg	45.0 (12.4)	11.5 (10.3)	74 (71%)	10.5 (17.4)	76 (73%)	50 (48%)	13 (13%)	11 (11%)	16 (15%)	32 (31%)
		Placebo	44.9 (12.8)	9.7 (7.7)	77 (73%)	7.9 (4.6)	75 (71%)	45 (43%)	13 (12%)	15 (14%)	11 (10%)	27 (26%)
SLEDAI-2 K	SLEDAI-2 K	SLEDAI-2 K organ system involvement	system involve	ement							PGA score, M	CLASI
score, M (SD)	score≥ 10, N (%)	CNS, N (%)	Vascular, N	Musculoskel-	Renal, N	Mucocutane-	Cardiovas-	Immunologi-	Constitu-	Hematolog-	(SD)	activity
			(0/2)	etal, N (%)	(06)	ous, N (%)	cular and respiratory,	Cal, N (%)	tional, N (%)	ICAI, N (%)		M (SD)
							N (%)					
10-1 (3-4)	149 (57%)	0	11 (4%)	253 (97%)	22 (8%)	252 (97%)	6 (2%)	143 (55%)	10 (4%)	19 (7%)	61-0 (12-5)	6.6 (6.6)
10.1 (3.0)	144 (56%)	0	14 (5%)	251 (97%)	10 (4%)	251 (97%)	9 (3%)	132 (51%)	11 (4%)	17 (7%)	58·8 (14·6)	6.7 (5.8)
10·1 (3·2)	147 (57%)	0	12 (5%)	252 (98%)	18 (7%)	243 (95%)	4 (2%)	138 (54%	8 (3%)	20 (8%)	60-0 (14-6)	6-8 (6-1)
10.3 (3)	152 (60%)	0	15 (6%)	250 (98%)	24 (9%)	246 (97%)	6 (2%)	131 (51%)	3 (1%)	13 (5%)	1-8 (0-4)	6·5 (7·0)
10-0 (3)	146 (58%)	0	10 (4%)	247 (98%)	17 (7%)	241 (96%)	8 (3%)	137 (54%)	2 (1%)	21 (8%)	1-8 (0-5)	6·1 (5·6)
10.1 (3)	144 (57%)	0	15 (6%)	247 (98%)	15 (6%)	244 (96%)	10 (4%)	134 (53%)	4 (2%)	13 (5%)	1-8 (0-5)	6-3 (6-2)
8-8 (3-4)	35 (33%)	1 (1%)	4 (4%)	93 (89%)	(%6) 6	82 (78%)	1 (1%)	63 (60%)	2 (2%)	6 (%6) (48·8 (15·8)	3.8 (5.4)
9.0 (3.3)	44 (42%	2 (2%)	3 (3%)	96 (92%)	7 (7%)	92 (88%)	1 (1%)	64 (62%)	0	5 (5%)	51.7 (16.0)	4.0 (3.4)
8.9 (2.9)	43 (41%)	3 (3%)	1 (1%)	93 (89%)	(%6) 6	90 (86%)	2 (2%)	62 (59%)	2 (2%)	13 (12%)	49.5 (16.9)	4.9 (5.7)
TJC, M (SD)	SJC, M (SD)	SLICC/ACR Dam-										
		age Index score, m (SD)										
10-3 (6-8)	6-64 (5-1)	0-68 (1-1)										
10.6 (7.1)	6.68 (5.0)	0.60 (1.0)										
10.6 (7.1)	6.41 (5.4)	0.66 (1.1)										
10.3 (6.5)	6.9 (5.1)	0.6 (1.0)										
10.9 (7.0)	6·9 (5·4)	0-6 (1-0)										
10.0 (7.0)	7.1 (5.6)	0-6 (1-0)										
8.7 (6.6)	5-2 (4-7)	0-44 (0-68)										
8·5 (6·2)	5-5 (4-2)	0-40 (0-88)										
7.7 (5.8)	5-3 (4-7)	0.59 (0.97)										

Diels Datia	




Fig. 4 A forest plot displaying the prevalence of Reduction of ≥4 points from baseline in Systemic Lupus Erythematosus Disease Activity Index 2000 score in the baricitinib group versus the placebo group

	Experim	ental	Contr	ol		Risk Ratio	Risk Ratio
Study or Subgroup	Events	Total	Events	Total	Weight	M-H, Fixed, 95% Cl	M-H, Fixed, 95% Cl
1.3.1 2 mg							
Morand 2023	126	255	121	253	19.7%	1.03 [0.86, 1.24]	+
Petri 2023	131	261	130	256	21.3%	0.99 [0.83, 1.17]	+
Wallace 2018	61	105	56	105	9.1%	1.09 [0.86, 1.39]	+
Subtotal (95% CI)		621		614	50.1%	1.02 [0.92, 1.14]	•
Total events	318		307				
Heterogeneity: Chi ² =	0.43, df = 1	2 (P = 0	.81); I² = I	0%			
Test for overall effect:	Z = 0.43 (F	^o = 0.67)				
1.3.2 4 mg							
Morand 2023	147	252	121	253	19.6%	1.22 [1.03, 1.44]	+
Petri 2023	133	258	130	256	21.2%	1.02 [0.86, 1.20]	+
Wallace 2018	70	104	56	105	9.1%	1.26 [1.01, 1.58]	
Subtotal (95% CI)		614		614	49.9%	1.14 [1.03, 1.27]	•
Total events	350		307				
Heterogeneity: Chi ² =	3.24, df = 3	2 (P = 0	20); I ^z = 3	38%			
Test for overall effect:	Z = 2.46 (F	^o = 0.01)				
Total (95% CI)		1235		1228	100.0%	1.08 [1.00, 1.17]	•
Total events	668		614				
Heterogeneity: Chi ² =	5.71, df = :	5 (P = 0	.34); I ² = 1	13%			0.01 0.1 1 10 100
Test for overall effect:	Z = 2.04 (F	^o = 0.04)				Favours [control] Favours [experimental]
Test for subgroup diffe	erences: C	≿hi² = 1.	92, df = 1	(P = 0.	.17), I² = 4	7.9%	ravours (control) - ravours (experimental)

Fig. 5 A forest plot displaying the prevalence of Systemic Lupus Erythematosus Disease Activity Index 2000 remission of arthritis or rash in the baricitinib group versus the placebo group

cebo group in terms of mild, moderate, or severe TEAEs. (Fig. 8)

than those in the placebo group [RR=1.49, 95% CI (1.16, 1.92), P=0.002]. (Figure S6)

Serious adverse events Participants in the baricitinib group were significantly more likely to obtain serious AEs

Discontinuation from study because of an adverse event Regarding Discontinuation from study treatment because of an AE, there was no significant differ-

	Experim	ental	Contr	ol		Risk Ratio		Risk Ratio	
Study or Subgroup	Events	Total	Events	Total	Weight	M-H, Fixed, 95% Cl		M-H, Fixed, 95% Cl	
1.4.1 2 mg									
Morand 2023	196	255	182	253	19.9%	1.07 [0.96, 1.18]		+	
Petri 2023	195	261	196	256	21.5%	0.98 [0.88, 1.08]		+	
Wallace 2018	82	105	80	105	8.7%	1.02 [0.88, 1.19]		+	
Subtotal (95% CI)		621		614	50.1%	1.02 [0.96, 1.09]		•	
Total events	473		458						
Heterogeneity: Chi ² =	= 1.58, df =	2 (P = 0	.45); l² = l	0%					
Test for overall effect	: Z = 0.64 (I	° = 0.52)						
1.4.2 4 mg									
Morand 2023	200	252	182	253	19.8%	1.10 [1.00, 1.22]		• •	
Petri 2023	188	258	196	256	21.4%	0.95 [0.86, 1.05]		-	
Wallace 2018	85	104	80	105	8.7%	1.07 [0.93, 1.23]		+	
Subtotal (95% CI)		614		614	49.9%	1.03 [0.97, 1.10]		•	
Total events	473		458						
Heterogeneity: Chi ² =	= 4.50, df =	2 (P = 0	.11); l ² = :	56%					
Test for overall effect	: Z = 1.00 (I	° = 0.32)						
Total (95% CI)		1235		1228	100.0%	1.03 [0.98, 1.07]			
Total events	946		916						
Heterogeneity: Chi2 =	= 6.18, df =	5 (P = 0	.29); I ² = 1	19%			L	0.1 1 10	1
Test for overall effect	: Z = 1.16 (I	^o = 0.25)				0.01	Favours [control] Favours [experim	
Test for subgroup dif	fferences: (chi² = 0.	06. df = 1	(P = 0)	.80), l ² = 0	1%		ravours (control) - ravours (experim	entaij






Fig. 6 A forest plot displaying the prevalence of No new BILAG A and no more than one new BILAB B disease activity score in the baricitinib group versus the placebo group. (A) before sensitivity analysis. (B) after sensitivity analysis

ence between the baricitinib group and the placebo group [RR=1.19, 95% CI (0.92, 1.54), P=0.19]. (Figure S7)

Infections and serious Infections The baricitinib group and the placebo group did not differ significantly in terms of infections [RR=1.03, 95% CI (0.95, 1.12), P=0.46].

However, participants in the baricitinib group were significantly more likely to obtain serious infections than those in the placebo group [RR=2.07, 95% CI (1.28, 3.34), P=0.003]. (Figures S8 and S9)

0 of 12




Fig. 7 A forest plot displaying the prevalence of No worsening by PGA in the baricitinib group versus the placebo group




Fig. 8 A forest plot displaying the prevalence of treatment emergent adverse events in the baricitinib group versus the placebo group

Opportunistic infection

Participants in the baricitinib group were significantly more likely to obtain opportunistic infection than those in the placebo group [RR=1.47, 95% CI (1.02, 2.11), P=0.04]. (Figure S10)

Hepatic disorders

The baricitinib group and the placebo group did not differ significantly in terms of hepatic disorders. (Figure S11)

Meta-regression

Random effect meta-regression was applied for dose of baricitinib, age, sample size of the study and time since

onset of SLE regarding SRI-4. There was no significant association between SRI-4 outcome and age of patients [OR=0.0104, 95% CI (-0.2488, 0.2697), P=0.937, figure S12], sample size [OR = -0.0005, 95% CI (-0.0024, 0.0014), P=0.6093, figure S13], dose of baricitinib [OR=0.0371, 95% CI (-0.2238, 0.2979), P=0.7805, figure S14], or time since onset of SLE[OR=0.1441, 95% CI (-0.1594, 0.4476), P=0.352, figure S15].

Discussion

Baricitinib has emerged as a potential therapeutic option for the treatment of wide scale autoimmune conditions including SLE due to its ability to modulate multiple cytokines involved in the pathogenesis of the disease [14]. Recent clinical trials have evaluated the safety and efficacy of baricitinib in SLE [15–17]. Herrin, we performed a meta-analysis utilizing data from these trials. Based on the literature search, we believe that our study is the first meta-analysis to evaluate the safety and efficacy of baricitinib in the treatment of SLE.

In terms of efficacy, the overall analysis demonstrated that baricitinib achieved the primary endpoint of a higher proportion of patients reaching an SRI-4 response compared with placebo. Our results were in line with the results of phase three RCT reported by Morand et al. [16]. In contrast, Petri et al. reported no significant SRI-4 response in baricitinib group compared with placebo [17]. The profile of patients, including organ involvement, in both trials was almost similar suggesting that patient clinical and immunological characteristics were not a major factor. Furthermore, the results of meta-regression analysis demonstrated no significant effect of age of patients, sample size, baricitinib dose, or time since onset on the primary outcome (SRI-4).

Furthermore, baricitinib demonstrated superiority over placebo in terms of BILAG A or B disease activity score. Additionally, it outperformed placebo in SLEDAI-2 K score of 4 points or more from baseline as well as a decrease in rash or arthritis based on the SLEDAI-2 K score. This adds a potential impact on reducing specific disease symptoms and activity. Our observations on these outcomes align with the results obtained from preclinical models. Preclinical studies conducted on (MRL/ lpr) mice and in vitro using immortalized primary podocytes and B cells isolated from C57BL/6 mice have demonstrated significant reductions in disease activity with baricitinib [22].

In terms of safety, Baricitinib appeared to be a well-tolerated drug for SLE. However, it is noteworthy that baricitinib had a higher incidence of serious adverse events and infections. This was similar to the results of pooled analysis on the safety of baricitinib in the treatment of SLE retorted by Dörner T et al. [23]. The results of our study should be interpreted cautiously because we were limited by the small number of clinical trials included. Only three randomized clinical trials were identified, which may have resulted in inaccurate precision of the results. Further research on the effectiveness of the baricitinib drug on patients with SLE is recommended. Moreover, long-term studies are also needed to assess the safety and efficacy of the drug over extended periods. Additionally, investigating the potential benefits of combining baricitinib with other treatments for SLE could provide valuable insights into the optimal management of this complex disease.

Conclusion

There is now sufficient clinical evidence to support Baricitinib's safety and efficacy in the treatment of SLE. This study's primary aim and some secondary endpoints were significantly reached by baricitinib. While glucocorticoids sparing and some other of the secondary outcomes didn't reach statistical significance, the overall analysis demonstrates a promising efficacy of baricitinib in the management of SLE.

List of abbreviations

BILAG	British Isles Lupus Assessment Group
LLDAS	Lupus low disease activity state
PGA	Physician Global Assessment
ROB	Risk of Bias Assessment tool
SLEDAI-2K	Systemic Lupus Erythematosus Disease Activity Index-2000)
SRI-4	Systemic Lupus Erythematosus Responder Index-4
TEAEs	Treatment-Emergent Adverse Events

Supplementary Information

The online version contains supplementary material available at https://doi.org/10.1186/s41927-023-00363-6.

Supplementary Material 1

Acknowledgements

Not applicable.

Authors' contributions

A.R.A. formulated and validated the research idea, prepared the data extraction sheets, revised the sheets and solved any conflicts, performed the meta-analysis and participated in writing and editing the final manuscript. M.S.A. performed data extraction, performed the quality assessment, revised the meta-analysis, and participated in writing and editing the final manuscript. A.M.M. validated the research idea performed data extraction, performed the quality assessment, revised the meta-analysis, participated in writing and editing the final manuscript.

Funding

The authors declare that no funds, grants, or other support was received during the preparation of this manuscript.

Data Availability

The datasets generated during and/or analyzed during the current study are available from the corresponding author upon reasonable request.

Declarations

Ethical approval

Not available.

Consent to publish Not applicable.

Competing interests

The authors declare no competing interests.

Received: 9 August 2023 / Accepted: 10 October 2023 Published online: 31 October 2023

References

- Tsokos GC, Lo MS, Costa Reis P, Sullivan KE. New insights into the immunopathogenesis of systemic Lupus Erythematosus. (1759–4804 (Electronic)).
- Tsokos GC. Systemic Lupus Erythematosus. N Engl J Med. 2011;365(22):2110–21.
- Fanouriakis A, Kostopoulou M, Alunno A, Aringer M, Bajema I, Boletis JN, et al. 2019 update of the EULAR recommendations for the management of systemic Lupus Erythematosus. Ann Rheum Dis. 2019;78(6):736–45.
- Kaul A, Gordon C, Crow MK, Touma Z, Urowitz MB, van Vollenhoven R, et al. Systemic Lupus Erythematosus. Nat Reviews Disease Primers. 2016;2:16039.
- Morand EF, Furie R, Tanaka Y, Bruce IN, Askanase AD, Richez C, et al. Trial of Anifrolumab in active systemic Lupus Erythematosus. N Engl J Med. 2020;382(3):211–21.
- Navarra SV, Guzman RM, Gallacher AE, Hall S, Levy RA, Jimenez RE, et al. Efficacy and safety of belimumab in patients with active systemic Lupus Erythematosus: a randomised, placebo-controlled, phase 3 trial. Lancet. 2011;377(9767):721–31.
- Wahren-Herlenius M, Dorner T. Immunopathogenic mechanisms of systemic autoimmune Disease. Lancet. 2013;382(9894):819–31.
- Larosa M, Zen M, Gatto M, Jesus D, Zanatta E, Iaccarino L, et al. IL-12 and IL-23/Th17 axis in systemic Lupus Erythematosus. Exp Biol Med (Maywood). 2019;244(1):42–51.
- Dorner T, Tanaka Y, Dow ER, Koch AE, Silk M, Ross Terres JA, et al. Mechanism of action of baricitinib and identification of biomarkers and key immune pathways in patients with active systemic Lupus Erythematosus. Ann Rheum Dis. 2022;81(9):1267–72.
- 10. Hoffman RW, Merrill JT, Alarcon-Riquelme MM, Petri M, Dow ER, Nantz E, et al. Gene expression and pharmacodynamic changes in 1,760 systemic

Lupus Erythematosus patients from two phase III trials of BAFF Blockade with Tabalumab. Arthritis Rheumatol. 2017;69(3):643–54.

- Schwartz DM, Kanno Y, Villarino A, Ward M, Gadina M, O'Shea JJ. JAK inhibition as a therapeutic strategy for immune and inflammatory Diseases. Nat Rev Drug Discovery. 2017;16(12):843–62.
- 12. Winthrop KL. The emerging safety profile of JAK inhibitors in rheumatic Disease. Nat Rev Rheumatol. 2017;13(4):234–43.
- Genovese MC, Kremer J, Zamani O, Ludivico C, Krogulec M, Xie L, et al. Baricitinib in patients with refractory rheumatoid arthritis. N Engl J Med. 2016;374:1243–52.
- King B, Ohyama M, Kwon OA-O, Zlotogorski A, Ko J, Mesinkovska NA et al. Two Phase 3 Trials of Baricitinib for Alopecia Areata. (1533–4406 (Electronic)).
- Wallace DJ, Furie RA, Tanaka Y, Kalunian KC, Mosca M, Petri MA, et al. Baricitinib for systemic Lupus Erythematosus: a double-blind, randomised, placebo-controlled, phase 2 trial. Lancet (London England). 2018;392(10143):222–31.
- Morand EF, Vital EM, Petri M, van Vollenhoven R, Wallace DJ, Mosca M et al. Baricitinib for systemic Lupus Erythematosus: a double-blind, randomised, placebo-controlled, phase 3 trial (SLE-BRAVE-I). Lancet (London, England). 2023.
- Petri M, Bruce IN, Dörner T, Tanaka Y, Morand EF, Kalunian KC et al. Baricitinib for systemic Lupus Erythematosus: a double-blind, randomised, placebocontrolled, phase 3 trial (SLE-BRAVE-II). Lancet. 2023.
- Tarsilla M. Cochrane handbook for systematic reviews of interventions. J Multidisciplinary Evaluation. 2010;6(14):142–8.
- 19. Takkouche B, Norman G. PRISMA statement. Epidemiology. 2011;22(1):128.
- Clark J, Glasziou P, Del Mar C, Bannach-Brown A, Stehlik P, Scott AM. A full systematic review was completed in 2 weeks using automation tools: a case study. J Clin Epidemiol. 2020;121:81–90.
- Sterne JA, Savović J, Page MJ, Elbers RG, Blencowe NS, Boutron I et al. RoB 2: a revised tool for assessing risk of bias in randomised trials. BMJ. 2019;366.
- 22. Lee J, Park Y, Jang SG, Hong SM, Song YS, Kim MJ et al. Baricitinib attenuates autoimmune phenotype and Podocyte Injury in a murine model of systemic Lupus Erythematosus. (1664–3224 (Electronic)).
- Dörner T, Tanaka Y, Mosca M, Bruce IN, Cardiel M, Morand EF, POS0714 pooled safety analysis of baricitinib in patients with systemic lupus erythematosus, et al. Results from three randomised, double-blind, placebo-controlled, clinical trials. Ann Rheum Dis. 2022;81 (Suppl 1):6391.

Publisher's Note

Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.